



## Health Literacy among Parents of Preschool Children with ADHD: A Cross-Sectional Study in Sanandaj, Iran

Nasrin Abdi<sup>1</sup>, Yaser Ghafouri<sup>2</sup>, Khaled Rahmani<sup>3</sup>, Ghazal Veisi<sup>4\*</sup>

1- Department of Community Health Nursing, Faculty of Nursing and Midwifery, Kurdistan University of Medical Sciences, Sanandaj, Iran.

2- General practitioner, Razi hospital, Baneh, Kurdistan university of medical Sciences, Sanandaj, Iran.

3- Associate Professor in Epidemiology, Liver and Digestive Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran.

4- Department of Community Health Nursing, School of Nursing and Midwifery, Kurdistan University of Medical Sciences, Sanandaj, Iran. **(Corresponding Author)** ghazallv98@gmail.com

**Background and Objectives:** ADHD is a chronic neurodevelopmental disorder affecting attention, activity levels, and impulse control, with a global prevalence of about 5% in children. In Iran, rates are similar, with the combined subtype being most common. ADHD impacts children's emotional and social development and places strain on families. Parental health literacy is essential for managing symptoms and improving outcomes; however, research on this topic especially among parents of preschool-aged children in Iran is limited. This study assessed the health literacy of parents whose preschool-aged children were diagnosed with attention-deficit/hyperactivity disorder (ADHD) and examined its association with ADHD in their children during 2022 in Sanandaj.

**Materials and Methods:** This descriptive-analytical study assessed parental health literacy using the "Health Literacy of Parents of Preschool Children" questionnaire, a researcher-developed instrument. Demographic information (including age, gender, education level, and employment status) was collected using a demographic questionnaire reviewed and approved by subject-matter experts. All parents of children diagnosed with ADHD were included through census sampling. Data were analyzed in Stata v.14 using independent t-tests and one-way ANOVA; non-parametric equivalents (Mann-Whitney U and Kruskal-Wallis tests) were applied when normality assumptions were unmet. A significance level of  $p < 0.05$  was considered.

**Results:** Among participants, 67.3% were female and 32.7% male, with a mean age of  $34.94 \pm 5.07$  years. The overall mean parental health literacy score was  $19.53 \pm 4.07$ . Nutrition subscale scores were significantly higher among parents of boys ( $p < 0.05$ ). Having a childcare book at home was significantly associated with both the developmental subscale and total literacy score ( $p < 0.05$ ). Parental education level showed significant relationships with nutrition, development, and numeracy subscales, as well as total literacy score ( $p < 0.05$ ). All subscales except development, along with total literacy score, were significantly associated with parental occupation ( $p < 0.05$ ).

**Conclusion:** Higher socioeconomic status and parental education were linked to increased health literacy. Owning childcare books and seeking accurate health information reflected greater parental responsibility and literacy.



**Keywords:** ADHD, Health Literacy, Parents, Attention Deficit Disorder with Hyperactivity, Children

**Received:** 02 November 2025

**Accepted:** 17 March 2026

**Doi:** 10.22038/jhl.2026.92417.1889

**Open Access Policy:** This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>

## Introduction

attention-deficit/hyperactivity disorder (ADHD) is a chronic brain disorder that manifests through ongoing issues with focus, excessive activity, and impulsiveness (1). ADHD impacts nearly 5% of children and about 2.5% of adults across the globe (2). Evidence from Iranian studies indicates that ADHD affects approximately 5 to 7 percent of children (3).

Three types of ADHD exist: inattentive, hyperactive-impulsive, and combined presentations. The last one is the most common subtype and it can disturb children's academic, social, and emotional development, often persisting into adulthood if left untreated (4).

The consequences of ADHD can deeply affect multiple aspects of life for both children and their parents. Its impact goes beyond the individual, placing emotional, relational, and financial pressure on families (1, 5).

Enhancing parental health literacy is a key determinant in preventing the exacerbation of symptoms and improving treatment outcomes in children with attention-deficit/hyperactivity disorder (ADHD). Parents with higher levels of health literacy are better able to understand basic information about the nature of ADHD, its natural course, and available treatment options (5). Moreover, interactive and critical health literacy aligned with Nutbeam's model empowers parents to communicate more effectively with healthcare providers, critically appraise health information, and engage more actively in shared decision-making regarding their child's care. Such informed engagement not only facilitates more effective healthcare utilization but also supports the implementation of parenting strategies such as Behavioral Parent Training (BPT) that align with therapeutic programs. This synergy, in turn, contributes to the reduction of ADHD symptoms and the enhancement of the child's overall functioning (6, 7).

Furthermore, parents with higher health literacy are better equipped to identify and mitigate environmental stressors such as irregular sleep patterns, poor nutrition, and family conflict thereby preventing the exacerbation of ADHD clinical presentations (8). In summary,



empowering parents through health literacy interventions enhances their knowledge base, psychological attitudes, and self-efficacy in symptom management. This, in turn, optimizes the ADHD therapeutic process and mitigates the long-term sequelae of the disorder.

According to the World Health Organization, health literacy reflects the personal knowledge and skills that are developed through daily activities, social interactions, and intergenerational transmission (9). Health literacy levels in Iran are low among the general population, similar to other developing countries (10). Recent research indicates that health literacy studies in early childhood tend to focus not on children themselves, but on the parents and educators who raise them highlighting that enhancing parental health literacy has a direct and positive impact on children's health status and behaviors (11). Nutbeam's Health Literacy Model is one of the most widely recognized and validated frameworks for understanding and promoting health literacy. This hierarchical model has been extensively utilized in school health education programs and public health interventions across diverse populations worldwide. It encompasses three progressive dimensions: functional, interactive, and critical health literacy. At the functional level, individuals develop essential reading, writing, and numeracy skills necessary to access and comprehend basic health information. The interactive level emphasizes effective communication, engagement with others, and the application of health information to modify behavior. At the critical level, individuals possess the ability to evaluate and reflect on health information, enabling them to make informed decisions regarding their own health and that of their communities. By highlighting individual empowerment and acknowledging environmental complexities, Nutbeam's model serves as a valuable tool for the design and evaluation of health promotion interventions (12). Parenting practices and the family environment play a significant role in shaping the emotional coping mechanisms of children with ADHD. Interventions that focus on parents have demonstrated greater effectiveness in reducing behavioral symptoms in affected children (13).

Available evidence suggests that parental health literacy in preschool-aged children has received limited research attention in Iran. This gap is evident both in terms of the domains required for comprehensive health literacy assessment and in the actual literacy levels among parents (14). Therefore, the present study aims to evaluate the health literacy of parents whose children have been diagnosed with attention-deficit/hyperactivity disorder (ADHD) and to examine its association with the presence of ADHD in their children in Sanandaj.

## **Materials and Methods**

This cross-sectional descriptive-analytical study was conducted to evaluate the health literacy levels of parents with children diagnosed with attention-deficit/hyperactivity disorder (ADHD).

### ***Participants***



The participants in this study included all parents of preschool children diagnosed with attention-deficit/hyperactivity disorder (ADHD), whose diagnosis had been confirmed through medical records. The contact information of parents was obtained by visiting pediatric psychiatric clinics, and they were subsequently contacted. The final sample size was 104 individuals.

## ***Procedures***

Following the approval of the Ethics Committee and written authorization from Kurdistan University of Medical Sciences, the study was conducted in coordination with academic supervisors. Contact information of parents of children diagnosed with attention-deficit/hyperactivity disorder (ADHD) was obtained by visiting pediatric psychiatric clinics. Parents were then contacted, informed about the study objectives, and invited to participate. Upon obtaining informed consent, data collection was carried out either through in-person visits coordinated with the treating physician or by sending the questionnaire via social media or email. Parental health literacy was assessed using the "Health Literacy of Parents of Preschool Children" questionnaire. Data were subsequently analyzed with the assistance of a statistical consultant, and the results were presented in the form of a thesis.

## ***Inclusion and Exclusion Criteria Inclusion Criteria***

According to the inclusion criteria, participants (either father or mother) had to be parents of preschool children diagnosed with ADHD, with the diagnosis confirmed by a psychiatrist and documented in the child's medical records. In addition, voluntary willingness to participate in the study was required.

In accordance with the exclusion criteria, participants who were unwilling to participate or chose to withdraw from the study at any stage were excluded.

## ***Data Collection***

Demographic information was completed by the parents using a researcher-designed questionnaire.

Demographic questions included parent gender (male/female), education level (less than high school diploma, diploma, bachelor's degree, master's degree or higher), and occupation (employed, unemployed, housewife), as well as the child's gender (male/female) and birth order (first, second, third or more) of the child diagnosed with ADHD. The questionnaire was developed based on expert input and underwent both face and content validation using qualitative methods. Face validity was assessed through feedback from 10 parents of preschool children, while content validity was confirmed by 12 experts in community health nursing and related fields.

Data were collected using the Health Literacy of Parents of Preschool Children questionnaire, which was distributed to parents either in person or via email.

Parental health literacy was assessed using the Health Literacy of Parents of Preschool Children questionnaire, which was researcher-developed and validated in the present study. The instrument consists of 40 items rated on a five-point Likert scale (1 = strongly disagree to 5 = strongly agree). The items are distributed across five domains: growth and development (15 items, score range 15–75), nutrition (9 items, score range 9–45), medical and health information (8 items, score range 8–40), numerical literacy (5 items, score range 5–25), and injury prevention (3 items, score range 3–15). Reliability of the instrument was confirmed through internal consistency (Kuder–Richardson coefficient = 0.78) and test–retest analysis (ICC = 0.81) (15).

### ***Data Analysis***

Data analysis in this study was performed using Stata software, version 14. Initially, descriptive statistics, including mean, standard deviation, frequency, percentage, and relevant charts, were used to summarize the data. To examine relationships between the study variables, appropriate statistical tests were applied based on the type of variable. The independent t-test was used to compare the mean health literacy scores across dichotomous variables, while one-way analysis of variance (ANOVA) was employed for variables with more than two categories. In cases where the distribution of the health literacy variable was not normal, equivalent non-parametric tests, namely the Mann–Whitney U test and the Kruskal–Wallis test, were used.

### ***Ethical Considerations***

Prior to enrollment, participants were fully informed about the purpose and procedures of the study. Written informed consent was obtained from all individuals before data collection. To ensure anonymity, no personal identifiers such as names or surnames were recorded on the questionnaires, which were coded numerically. All participant data were treated as confidential and securely maintained by the researcher.

## **Results**

### ***Sociodemographic Characteristics of Parents***

Among the 104 participating parents, the majority were female (67.3%) and housewives (59.6%). The mean age was 34.94 years (SD = 5.07), ranging from 24 to 47 years. Most had one child (55.8%), and in 75% of cases, the child with ADHD was the first-born. More than half of the parents (54.8%) had less than a high school diploma, while only 6.7% held a master's degree or higher. Most had not attended any educational classes related to child health (88.5%), although 91.3% reported access to pediatric care books. Regarding sources of childcare information, the most common were consulting physicians or health centers (47.1%) and using the internet (32.7%). Broadcast media exposure was limited, with 71.2%



reporting no use of such resources. In terms of household income, 35.6% of families earned more than 600,000 Tomans per month, while 19.2% earned less than 250,000 Tomans (Table 1).

**Descriptive Analysis Parental Health Literacy and its Dimensions**

The mean total health literacy score among participating parents was 19.53 (SD = 4.84), with scores ranging from 8 to 31. Among the dimensions, the highest mean score was observed in the growth and development domain (6.26 ± 2.23), followed by nutrition (4.84 ± 1.84) and medical and health information (4.50 ± 1.28). Lower scores were reported in numeracy (2.16 ± 1.46) and safety and injury prevention (1.77 ± 0.84). These findings suggest that while parents demonstrated moderate literacy in developmental and nutritional aspects, their numeracy and safety-related literacy were relatively limited (Table 2).

**Table 1. Sociodemographic Characteristics of Parents**

Variables	Values
<b>Gender</b>	
Male	34 (32.7)
Female	70 (67.3)
Total	104 (100.0)
<b>Age</b>	
Minimum	24
Maximum	47
Mean ± SD	34.94 ± 5.07
<b>Number of Children</b>	
1	58 (55.8)
2	34 (32.7)
3	10 (9.6)
5	2 (1.9)
<b>ADHD Child's Birth Order</b>	
1	78 (75.0)
2	20 (19.2)
3	5 (4.8)
4	1 (1.0)
<b>Job</b>	
Housewife	62 (59.6)
Employee	29 (27.9)
Other	13 (12.5)
<b>Participation in Educational Classes</b>	
No participation	92 (88.5)
Less than one week	9 (8.7)
More than one week	3 (2.9)
<b>Education</b>	



Less than high school diploma	57 (54.8)
Associate degree	19 (18.3)
Bachelor's degree	21 (20.2)
Master's degree or higher	7 (6.7)
<b>Access to Pediatric Care Books</b>	
Yes	95 (91.3)
No	9 (8.7)
<b>Reasons for Unavailability of Childcare Books</b>	
Negligence	45 (47.4)
Lack of time	29 (30.5)
No perceived need	15 (15.8)
cost of books	3 (3.2)
Lack of awareness	3 (3.2)
<b>Exposure to Health Education via Broadcast Media</b>	
Not used	74 (71.2)
Television	15 (14.4)
Radio	3 (2.9)
Social media	12 (11.5)
<b>Sources of Childcare Information Used by Parents</b>	
Consulting elders and relatives	10 (9.6)
Using books	11 (10.6)
Using the internet	34 (32.7)
Consulting physicians or health centers	49 (47.1)
<b>Monthly Household Income (×1,000 Toman)</b>	
Less than 250	20 (19.2)
250 to 400	30 (28.8)
400 to 600	17 (16.3)
More than 600	37 (35.6)

a Variables are expressed as No. (%) or mean ± SD (Min - Max)

**Table 2. Descriptive Analysis Parental Health Literacy and its Dimensions (n = 104)**

Subscale	Minimum	Maximum	Mean ± SD
Nutrition	0.00	9.00	4.84 ± 1.84
Growth and Development	0.00	11.00	6.26 ± 2.23
Safety and Injury Prevention	0.00	3.00	1.77 ± 0.84
Numeracy	0.00	5.00	2.16 ± 1.46
Medical and Health Information	1.00	7.00	4.50 ± 1.28
<b>Total Health Literacy Score</b>	8.00	31.00	19.53 ± 4.84

**Association Between Sociodemographic Variables and Health Literacy Dimensions**

Based on Table 3, the association between sociodemographic variables and dimensions of health literacy is presented.

Female parents scored significantly higher in the nutrition domain compared to males (5.56 ± 1.71 vs. 4.49 ± 1.80, p = 0.005). No significant differences were observed in other

dimensions ( $p > 0.05$ ). No statistically significant differences were found across any health literacy domains ( $p > 0.05$ ), although numeracy scores were notably lower among parents with five children ( $0.00 \pm 0.00$ ). Birth order was not significantly associated with any health literacy dimensions ( $p > 0.05$ ). Significant differences were observed in all domains except growth and development. Employed parents had higher scores in nutrition ( $p = 0.007$ ), safety ( $p = 0.009$ ), numeracy ( $p = 0.005$ ), medical information ( $p = 0.010$ ), and total health literacy ( $p = 0.0001$ ).

No significant associations were found ( $p > 0.05$ ), although parents who attended classes for more than one week had the highest scores in growth and development ( $9.00 \pm 1.73$ ,  $p = 0.076$ ). Statistically significant differences were found in nutrition ( $p = 0.001$ ), growth and development ( $p = 0.005$ ), numeracy ( $p = 0.033$ ), and total health literacy ( $p = 0.001$ ). Parents with a master's degree or higher consistently scored highest across these domains. Parents with access to pediatric books had significantly higher scores in growth and development ( $p = 0.014$ ) and total health literacy ( $p = 0.042$ ). Significant differences were found in medical information ( $p = 0.049$ ) and total health literacy ( $p = 0.003$ ). Parents who used social media had the highest scores across most domains. Significant associations were observed in nutrition ( $p = 0.029$ ), growth and development ( $p = 0.020$ ), numeracy ( $p = 0.035$ ), and total health literacy ( $p = 0.001$ ). Parents who relied on the internet or books scored higher than those who consulted elders or health centers.

**Table 3. Association between Demographic Variables and Dimensions of Health Literacy**

Demographic Variable	Nutrition	Growth & Development	Safety	Numeracy	Medical Info	Total Score
<b>Gender</b>						
Male	4.49 ± 1.80	6.21 ± 2.48	1.74 ± 0.85	2.13 ± 1.36	4.40 ± 1.15	18.97 ± 4.77
Female	5.56 ± 1.71	6.35 ± 1.63	1.82 ± 0.83	2.24 ± 1.65	4.71 ± 1.51	20.68 ± 4.85
<b>p-value</b>	<b>0.005</b>	<b>0.768</b>	<b>0.648</b>	<b>0.728</b>	<b>0.254</b>	<b>0.096</b>
<b>Number of Children</b>						
1 child	5.05 ± 1.67	6.52 ± 2.04	1.81 ± 0.83	2.17 ± 1.48	4.41 ± 1.35	19.96 ± 4.09
2 children	4.65 ± 2.09	5.68 ± 2.50	1.76 ± 0.89	2.44 ± 1.37	4.53 ± 1.28	19.06 ± 5.73
3 children	4.60 ± 1.96	6.80 ± 2.44	1.70 ± 0.82	1.60 ± 1.35	5.00 ± 0.82	19.70 ± 5.77
5 children	3.50 ± 0.71	6.00 ± 0.00	1.00 ± 0.00	0.00 ± 0.00	4.00 ± 0.00	14.50 ± 0.71
<b>p-value</b>	<b>0.504</b>	<b>0.299</b>	<b>0.602</b>	<b>0.065</b>	<b>0.554</b>	<b>0.403</b>
<b>ADHD Child's Birth Order</b>						
First-born	5.01 ± 1.87	6.21 ± 2.32	1.73 ± 0.86	2.18 ± 1.47	4.42 ± 1.31	19.55 ± 4.98
Second-born	4.40 ± 1.82	6.45 ± 1.70	1.95 ± 0.69	2.50 ± 1.32	4.80 ± 1.28	20.10 ± 4.04
Third-born	4.20 ± 1.30	6.40 ± 3.36	1.80 ± 1.10	1.00 ± 1.22	4.60 ± 0.55	18.00 ± 6.20
Fourth-born	4.00 ± -	6.00 ± -	1.00 ± -	0.00 ± -	4.00 ± -	15.00 ± -
<b>p-value</b>	<b>0.454</b>	<b>0.974</b>	<b>0.591</b>	<b>0.088</b>	<b>0.672</b>	<b>0.653</b>
<b>Job</b>						
Housewife	4.44 ± 1.88	6.02 ± 2.41	1.74 ± 0.79	2.05 ± 1.32	4.39 ± 1.15	18.62 ± 4.74
Employee	5.72 ± 1.69	6.83 ± 1.98	2.07 ± 0.88	2.79 ± 1.50	5.03 ± 1.24	22.45 ± 3.97
Other	4.77 ± 1.30	6.15 ± 1.77	1.23 ± 0.73	1.31 ± 1.49	3.85 ± 1.57	17.31 ± 4.46
<b>p-value</b>	<b>0.007*</b>	<b>0.269</b>	<b>0.009*</b>	<b>0.005*</b>	<b>0.010*</b>	<b>0.0001*</b>



Participation in Educational Classes						
None	4.76 ± 1.91	6.13 ± 2.19	1.76 ± 0.84	2.09 ± 1.48	4.52 ± 1.30	19.25 ± 4.91
< 1 week	5.78 ± 0.67	6.67 ± 2.40	1.89 ± 0.78	2.67 ± 0.87	4.44 ± 1.01	21.44 ± 3.91
> 1 week	4.67 ± 1.15	9.00 ± 1.73	1.67 ± 1.15	3.00 ± 2.00	4.00 ± 1.73	22.33 ± 4.04
<b>p-value</b>	<b>0.281</b>	<b>0.076</b>	<b>0.890</b>	<b>0.316</b>	<b>0.781</b>	<b>0.260</b>
Education						
Less than diploma	4.30 ± 1.94	5.86 ± 2.20	1.67 ± 0.83	1.79 ± 1.45	4.32 ± 1.24	17.91 ± 5.00
Associate degree	6.11 ± 1.15	6.00 ± 2.05	1.68 ± 0.82	2.53 ± 1.26	4.37 ± 1.50	20.68 ± 3.84
Bachelor's degree	5.24 ± 1.61	6.71 ± 2.19	1.90 ± 0.83	2.62 ± 1.32	5.00 ± 1.05	21.48 ± 4.24
Master's and above	4.57 ± 1.27	8.86 ± 1.21	2.43 ± 0.79	2.86 ± 1.77	4.86 ± 1.35	23.57 ± 2.07
<b>p-value</b>	<b>0.001*</b>	<b>0.005*</b>	<b>0.114</b>	<b>0.033*</b>	<b>0.161</b>	<b>0.001*</b>
Access to Pediatric Care Books						
Yes	4.67 ± 1.00	8.00 ± 1.58	2.22 ± 0.83	3.00 ± 1.50	4.78 ± 0.67	22.67 ± 2.92
No	4.86 ± 1.90	6.09 ± 2.22	1.73 ± 0.83	2.08 ± 1.43	4.47 ± 1.32	19.23 ± 4.89
<b>p-value</b>	<b>0.762</b>	<b>0.014*</b>	<b>0.090</b>	<b>0.071</b>	<b>0.497</b>	<b>0.042*</b>
Exposure to Health Education via Broadcast Media						
No use	4.62 ± 1.93	6.07 ± 2.31	1.80 ± 0.84	2.01 ± 1.34	4.61 ± 1.20	19.10 ± 4.79
Television	5.47 ± 0.83	6.93 ± 1.75	1.73 ± 0.59	2.60 ± 1.59	4.20 ± 1.61	20.93 ± 3.77
Radio	3.67 ± 2.31	4.00 ± 2.65	0.67 ± 0.58	1.00 ± 1.73	2.67 ± 0.58	12.00 ± 2.65
Social media	5.75 ± 1.66	7.17 ± 1.70	1.92 ± 1.00	2.83 ± 1.70	4.67 ± 1.07	22.33 ± 4.42
<b>p-value</b>	<b>0.069</b>	<b>0.071</b>	<b>0.128</b>	<b>0.084</b>	<b>0.049*</b>	<b>0.003*</b>
Sources of Childcare Information Used by Parents						
Elders and relatives	5.30 ± 1.57	6.60 ± 1.84	1.80 ± 0.63	1.80 ± 1.14	4.50 ± 1.08	5.30 ± 1.57
Books	5.27 ± 1.95	6.27 ± 2.15	1.91 ± 0.70	2.91 ± 1.14	4.45 ± 1.37	5.27 ± 1.95
Internet	5.38 ± 1.35	7.12 ± 1.93	1.88 ± 0.81	2.53 ± 1.46	4.82 ± 1.36	5.38 ± 1.35
Physicians / Health centers	4.27 ± 2.03	5.59 ± 2.35	1.65 ± 0.93	1.82 ± 1.48	4.29 ± 1.22	4.27 ± 2.03
<b>p-value</b>	<b>0.029*</b>	<b>0.020*</b>	<b>0.604</b>	<b>0.035*</b>	<b>0.314</b>	<b>0.001*</b>

## Discussion

The findings of this descriptive-analytical study expand the current understanding of factors associated with parental health literacy in families of preschool children with ADHD. These results can be interpreted in the context of previous studies, while taking into account the specific cultural background and study population.

Overall, the findings of this study showing a positive association between parental education level, occupational status, and access to structured information resources (such as educational books) with health literacy scores are consistent with a broad body of similar research. For example, in line with the study by Olsbø et al. (2024) on parents of children with Hirschsprung disease, higher educational attainment was identified as a strong determinant of health literacy. Likewise, although the study by Valaei Sharif et al. (2025) focused on the effectiveness of an educational intervention, it implicitly confirmed the importance of parents' baseline knowledge. In addition, Dekkers et al. (2020) demonstrated that parental knowledge mediated the relationship between ADHD symptoms and behavioral outcomes in adolescents, highlighting the broader implications of parental literacy for child development.



Similarly, the findings of Csima (2024) and Uner Asil (2025) indicated that parental health literacy was associated with healthier habits and better overall health status in children. Taken together, these studies align with the present findings. The distinctive contribution of the current study lies in its focus on parents of preschool children with ADHD in Sanandaj, a population facing unique caregiving challenges and potential limitations in social and educational infrastructures (10, 16-19).

The primary aim of this study was to assess parental health literacy and identify associated factors. In this regard, the strong association observed between parental education level, socioeconomic status, and health literacy should not be interpreted as causal, yet it remains noteworthy. Higher educational attainment likely strengthens basic functional literacy skills (such as reading and comprehension) as well as critical literacy skills (such as evaluating information sources). Moreover, better occupational and economic status often translates into greater access to high-quality educational resources and the time and psychological capacity to actively seek information. Taken together, these factors may create a cycle in which social inequalities contribute to disparities in health literacy and, consequently, in the quality of child care.

Another noteworthy finding of our study was the stronger association of digital and printed resources, compared to consulting family elders, with higher parental health literacy. This result is consistent with the study conducted in Saku, Japan (2022), which showed that access to structured content through a mobile application enhanced parents' child-care knowledge and health literacy (20). This similarity suggests that the quality and structured nature of information, regardless of whether it is delivered digitally or in print, is a key factor. However, differences may lie in the level of access to and trust in these resources across different populations.

access to child-care books, as a tangible indicator of an information-enriched environment, was associated with higher parental health literacy. This finding suggests that merely emphasizing individual knowledge acquisition is insufficient; rather, ensuring equitable and easy access to reliable, clear, and culturally appropriate information resources is essential particularly for parents who may face linguistic or educational barriers in searching for or evaluating information. This underscores the importance of policy initiatives at the macro level (such as the production and distribution of standardized resources) alongside the design of targeted interventions at the micro level

the study by Goodwin et al. (2023) on children with medical complexity did not find a significant association between low parental health literacy and the frequency of emergency visits, but reported substantially higher care costs. This suggests that the consequences of inadequate health literacy may not always manifest as increased service utilization, but rather as poorer quality of disease management and, ultimately, greater financial burden (21).



Despite differences in the study population, this finding indirectly underscores the importance of parental health literacy for the efficient and cost-effective management of chronic conditions such as ADHD.

Parental socioeconomic status was found to be associated with health literacy. This result is consistent with the studies of Csima (2024) and Uner Asil (2025), which demonstrated that parental health literacy was linked to healthier habits and better overall health outcomes in children. The differences observed across studies may be explained by variations in access to educational resources, differences in social support systems, and the level of cultural capital within families. Taken together, these findings highlight the importance of supportive policies aimed at reducing social inequalities and improving access to educational resources so that parents from diverse socioeconomic backgrounds have equal opportunities to enhance their health literacy (18).

**Study Limitations and Strengths:** The interpretation of this study's findings should take into account several limitations. First, the cross-sectional design precludes causal inference; that is, it cannot be determined whether higher education directly leads to better health literacy or whether other variables play a role. Second, the sampling method, restricted to a single city (Sanandaj) with a relatively small sample size, may limit the generalizability of the results to other regions of Iran with different cultural and socioeconomic contexts. Third, reliance on parents' self-reported data for assessing health literacy and behaviors may have been influenced by social desirability bias. Fourth, although some sociodemographic factors were controlled, important variables such as parental stress level, social support, and the severity of the child's symptoms were not measured or adjusted for, which could have affected the observed associations. Finally, although the questionnaire used demonstrated acceptable validity and reliability, it was a general tool for parental health literacy and may not have captured specific dimensions relevant to ADHD management, such as mental health literacy or medication literacy.

## Conclusion

This study indicates that among parents of preschool children with ADHD in Sanandaj, factors such as educational level, occupational status, and access to structured information resources are associated with higher levels of health literacy. Although these associations are not causal in nature, they highlight critical points for future interventions. Effective educational programs for these parents should be tailored to their baseline literacy level, cultural context, and available resources, with a focus on facilitating access to reliable and practical information. Future research employing longitudinal designs and ADHD-specific assessment tools could provide a deeper understanding of the dynamic relationship between parental health literacy and children's long-term outcomes.



Mashhad University of Medical Sciences



**Acknowledgements:** We would like to express our sincere gratitude to all individuals who contributed to the completion of this study.

**Availability of Data and Materials:** The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

**Conflicts of interest:** The authors state that there is no conflict of interest in the present study.

**Consent for publication:** Not applicable.

**Ethical Approval and consent to participate:** Ethical approval for this study was granted by the Ethics Committee of Kurdistan University of Medical Sciences under the code IR.MUK.REC.1401.079, and informed consent was obtained based on the Deceleration of Helsinki.

**Funding:** This study received financial support from the Vice President of Research and Technology, Kurdistan University of Medical Sciences.

**Author Contributions:** N.A. Y.GH., KH.R., and GH.V. contributed to the study design, data collection, and initial manuscript drafting. Statistical analysis was performed by KH.R., N.A. supervised the study design, implementation, and manuscript writing. All authors read and approved the final version of the manuscript.

## References

1. Shafighi A, Mousavizadeh S, Koshki M, Fini Y. Health Literacy and Parenting Stress Among Iranian Parents of Children with Attention Deficit Hyperactivity Disorder: A Cross-Sectional Study. *Middle East Journal of Rehabilitation and Health Studies*. 2024;11.
2. Paiva GCC, de Paula JJ, Costa DS, Alvim-Soares A, Santos D, Jales JS, et al. Parent training for disruptive behavior symptoms in attention deficit hyperactivity disorder: a randomized clinical trial. *Front Psychol*. 2024;15:1293244.
3. Tajik A, Nikfar S, Elyasi S, Rajabi O, Varmaghani M. Cost-effectiveness and budget impact analysis of lisdexamfetamine versus methylphenidate for patients under 18 with attention-deficit/hyperactivity disorder in Iran. *Child Adolesc Psychiatry Ment Health*. 2023;17(1):115.
4. Patel P, Behl J, Karim T, Palnati SR, Bhakta S. Parenting Styles and Attention Deficit Hyperactivity Disorder in the South Asian Population: A Narrative Review. *Cureus*. 2024;16(12):e74967.
5. Sharif NV, Ghasemzadeh P, Mohebbi N, Ghasemzadeh S. Improved medication adherence and health literacy in parents of children with ADHD: Effects of a targeted educational program. *Explor Res Clin Soc Pharm*. 2025;19:100634.
6. Breaux R, Waschbusch DA, Marshall R, Humphrey H, Pelham WE, Jr., Waxmonsky JG. The Role of Parental Knowledge and Attitudes about ADHD and Perceptions of Treatment Response in the Treatment Utilization of Families of Children with ADHD. *Evid Based Pract Child Adolesc Ment Health*. 2020;5(1):102-14.
7. Li D, Feng X, Yuan Y, Li Y, Zhang L, Tian T, et al. Behavioral parent training interventions to improve health outcomes of preschool children with ADHD symptoms from Chinese disadvantaged families: study protocol for a randomized controlled trial. *Trials*. 2025;26(1):497.
8. Brown L, Tallon M, Bellgrove M, Rudaizky D, Kendall G, Boyes M, et al. Increasing Health Literacy on ADHD: A Cross-Disciplinary Integrative Review Examining the Impact of ADHD on Brain Maturation, Composition and Function and Cognitive Processes Across the Life Course. *Child Psychiatry & Human Development*. 2025:1-39.



Mashhad University of Medical Sciences



9. Organization WH. Health literacy – Fact sheet: World Health Organization; 2024 [Available from: <https://www.who.int/news-room/fact-sheets/detail/health-literacy>].
10. Sharif NV, Ghasemzadeh P, Mohebbi N, Ghasemzadeh S. Improved medication adherence and health literacy in parents of children with ADHD: Effects of a targeted educational program. *Exploratory Research in Clinical and Social Pharmacy*. 2025;100634.
11. Feng M, Xu J, Zhai M, Wu Q, Chu K, Xie L, et al. Behavior Management Training for Parents of Children with Preschool ADHD Based on Parent-Child Interactions: A Multicenter Randomized Controlled, Follow-Up Study. *Behav Neurol*. 2023;2023:3735634.
12. Peralta LR, Forsyth R, Lapi N, Amon KL, Metsan P, Wattlez G, et al. Health literacy of adolescents' responses to a workshop focusing on food, nutrition, climate change and digital technology solutions in Oceania: a multi-site pilot study in Vanuatu. *BMC Public Health*. 2025;25(1):648.
13. Bringer M, Bodard S, Moscoso A, Revah-Levy A, Purper-Ouakil D, Acquaviva E, et al. Online search and activities of parents of children with ADHD: a qualitative study. *Child Adolesc Psychiatry Ment Health*. 2025;19(1):31.
14. Mozafari M, Borji M. Assessing the health literacy level of parents with school children in Ilam, 2015. *Journal of Nursing Education*. 2017;5(6):53-61.
15. Chehri M. Assessment of Health Literacy Levels of Parents of Preschool Children. *Journal of Pediatric and Neonatal Nursing*. 2015; 1 (4):1–10.
16. Olsbø S, Kiserud SG, Hermansen Å, Hamilton Larsen M, Bjørnland K. Health literacy in parents of children with Hirschsprung disease: a novel study. *Pediatr Surg Int*. 2024;41(1):21.
17. Dekkers TJ, Huizenga HM, Bult J, Popma A, Boyer BE. The importance of parental knowledge in the association between ADHD symptomatology and related domains of impairment. *Eur Child Adolesc Psychiatry*. 2021;30(4):657-69.
18. Uner Asil E, Aygun O, Duman A, Gungor N. The relationship between parental health literacy and primary school students' anthropometric measurements and general health status. *BMC Pediatr*. 2025;25(1):125.
19. Csimá M, Podráczky J, Keresztes V, Soós E, Fináncz J. The Role of Parental Health Literacy in Establishing Health-Promoting Habits in Early Childhood. *Children (Basel)*. 2024;11(5).
20. Sakamoto M, Ishikawa H, Suzuki A. Evaluation of Parents' Use of a Child Health Care Information App and Their Health Literacy: Cross-Sectional Study. *JMIR Pediatr Parent*. 2024;7:e48478.
21. Goodwin EJ, Zaniletti I, Solano J, Bettenhausen JL, Coller RJ, Plencner LM, et al. Parental Health Literacy and Acute Care Utilization in Children With Medical Complexity. *Hosp Pediatr*. 2024;14(10):e426-e31.